

FAUNA VILINIH KONJICA

Bosne i Hercegovine



Calopteryx splendens (Harris, 1782)



Calopteryx virgo (Linnaeus, 1758)



Lestes virens (Charpentier, 1825)



Lestes viridis (Vander Linden, 1825)



Sympecma fusca (Vander Linden, 1820)



Enallagma cyathigerum Charpentier, 1840



Erythromma viridulum (Charpentier, 1840)



Ischnura elegans (Vander Linden, 1820)



Coenagrion puella (Linnaeus, 1758)



Pyrrhosoma nymphula (Sulzer, 1776)



Ceriagrion tenellum (de Villers, 1789)



Platycnemis pennipes (Pallas, 1771)



Callaeschna microstigma (Schneider, 1845)



Gomphus vulgatissimus (Linnaeus, 1758)



Cordulegaster heros Theischinger, 1979



Cordulegaster bidentata Selys, 1843



Somatochlora flavomaculata
(Vander Linden, 1825)



Libellula quadrimaculata Linnaeus, 1758



Orthetrum albistylum (Selys, 1848)



Orthetrum coerulescens (Fabricius, 1798)



Sympetrum sanguineum (Müller, 1764)



Sympetrum meridionale (Selys, 1841)



Crocothemis erythraea (Brullé, 1832)

Vilini konjici, insekti iz reda Odonata, spadaju među najveće i najspektakularnije insekte koji, gotovo nepromijenjeni, postoje na Zemlji već više od 300 miliona godina. Najveći dio života ovi insekti provode u stadiju larve skriveni ispod površine vode, dok su kao odrasli vezani za različite tipove kopnenih staništa. Oni su svojevrsni čuvari prirode, a njihovo prisustvo pokazatelj je bogatstva njihovih staništa. Vilini konjici među prvima reaguju na promjene u životnoj sredini i zbog toga su važan indikator stanja i očuvanosti različitih tipova slatkovodnih ekosistema.

Od oko 140 vrsta vilinih konjica poznatih u Evropi, 15% je ugroženo, 11% je gotovo ugroženo, a 24% bilježi pad brojnosti. U Bosni i Hercegovini su do sada registrovane 63 vrste vilinih konjica među kojima su dvije vrste ugrožene u Evropi kao i četiri vrste koje su zaštićene prema Evropskoj Direktivi o vrstama i staništima. U Evropi i Svijetu već dugo vremena je poznat značaj i ugroženost ovih insekata zbog čega su brojne vrste i njihova staništa stavljeni na listu zaštićenih, međutim u Bosni i Hercegovini niti jedna vrsta vilinih konjica još uvijek nije zaštićena, a njihova staništa su svakodnevno izložena uništavanju.

Dragonflies of Hutovo Blato



*Calopteryx splendens
balcanica* Fudakowski, 1930



Calopteryx virgo
(Linnaeus, 1758)



Sympecma fusca
(Vander Linden, 1820)



Enallagma cyathigerum
Charpentier, 1840



Anax ephippiger (Burmeister, 1839)



Coenagrion puella
(Linnaeus, 1758)



Erythromma viridulum
(Charpentier, 1840)



Ceragrion tenellum
(de Villers, 1789)



Platycnemis pennipes
(Pallas, 1771)



Gomphus vulgatissimus (Linnaeus, 1758)



Aeshna affinis
Vander Linden, 1820



Brachytron pratense
(Müller, 1764)



Onychogomphus forcipatus
(Linnaeus, 1758)



Libellula depressa
Linnaeus, 1758



Libellula fulva
(Müller, 1764)



Orthetrum cancellatum
(Linnaeus, 1758)



Orthetrum coerulescens
(Fabricius, 1798)



Crocothemis erythraea
(Brullé, 1832)



Sympetrum sanguineum (Müller, 1764)